


Protein Marker PS10 PLUS

Description/Preparation:

The Protein Marker PS10 is a mixture of 10 purified proteins which are re-dissolved 'ready-to-use' in loading buffer. The proteins resolve into 10 sharp bands in the range of 11-180 kDa when analysed by SDS-PAGE and stained with optimized dyes. **The marker is ready-to-use. There is no need to boil.**



Applications

- Monitoring of protein migration during SDSpolyacrylamide gel electrophoresis.
- Monitoring of protein transfer onto membranes during Western blotting.
- Sizing of proteins on SDS-polyacrylamide gels and Western blots.

Usage:

Mini gel application: 10 μ l/well; 5 μ l per well blots Standard gel application: 20 μ l/well; 10 μ l per well for blots

Number of bands: 10 11, 17, 25, 35, 48, 63, 75, 100.135, 180 kDa

Loading:

Loading Denaturing Polyacrylamide gels (SDS-PAGE):

- Thaw marker at room temperature or heat at 37 40 °C for a few minutes. Do not boil!
- Vortex gently to ensure the solution is homogeneous and load the ladder on SDSpolyacrylamide gel
- 10 μl per well for mini-gels, 5 μl per well for blots
- 20 µl per well for large gels, 10 µl per well for blots

1 ml marker is sufficient for 100 mini gels or 50 standard gels.

It is recommended to divide the marker into aliquots to avoid contamination of the stock solution.

Note:

Protein Marker PS10 is optimized for runs on 12 % SDS polyacrylamide gels. 8 to 10 % gels may cause proteins with low molecular weights to migrate with the dye front. On 12 to 15 % and gradient gels all bands are visible.

Quality Control

Tested in SDS-polyacrylamide gel electrophoresis and Western blotting.

Storage: at -20°C

Shipment: on blue ice

Ordering Information:

CatNo.	Description	Amount
310003	Protein Marker PS10 PLUS (11-180 kDa)	500 µl
310004	Protein Marker PS10 PLUS (11-180 kDa)	5 x 500 μl

μΙ ο μΙ decision.