

Ribonuklease Inhibitor (RNase Inhibitor)

Applications for RNase inhibitor:

- Inhibits common eukaryotic RNases
- Active over a broad pH range (pH 5-8)
- high levels of inhibition over a wide range of conditions

Description:

RNase Inhibitor is a recombinant human placental protein which inhibits ribonucleases (RNases) A, B and C. It does not inhibit RNase 1, RNase T1, S1 Nuclease, RNase H or RNase from *Aspergillus*. There is no inhibition of polymerase activity when the protein is used with Taq DNA Polymerase, AMV or M-MuLV Reverse Transcriptases, or Phage RNA Polymerases (SP6, T7, or T3).

Concentration: 40 u/μl

Storage Buffer:

20 mM HEPES-KOH (pH7.6), 50 mM KCl, 8 mM DTT and 50% glycerol

Unit definition RNase inhibitor:

One unit is the amount of enzyme required to inhibit by 50% the activity of 5 ng of RNase A at 25°C (This inhibitor activity is determined by its ability to inhibit hydrolysis of cyclic 2', 3'-CMP by RNase A).

Quality Control: Endonuclease: Contains no detectable endonuclease activity. Incubation of 200 units of enzyme with supercoiled plasmid produced no nicked molecules after a two hour incubation at 37°C as determined by ethidium-stained agarose gel electrophoresis.

Ribonuclease: No ribonuclease activity is observed after 1 μg of RNA is incubated with 200 units of enzyme for 60 minutes at 37°C. The RNA is electrophoresed on an agarose gel and stained with ethidium bromide. No latent ribonuclease activity is observed after 1 μg of RNA is incubated with 200 units of pre-heated enzyme for 60 minutes at 37°C. The RNA is electrophoresed on an agarose gel and stained with ethidium bromide.

DNase: 50 ng of radiolabelled DNA is incubated with 200 units of enzyme for 60 minutes at 37°C, and the release of radiolabelled nucleotides is monitored by scintillation counting of TCA-soluble material. Minimum passing specification is <3% release of input radioactivity into TCA-soluble material.

Usage or RNase inhibitor:

We recommend to use 1 unit per reaction unit

Note:

- Ribonuclease Inhibitor requires at least 1 mM DTT to be active,
- Enzyme inhibits in a wide pH range, but most strongly at pH 7 - 8
- avoid temperatures above 50 °C and high concentrations of urea or other denaturing agents

Transportation: RNase inhibitor is shipped on blue ice

Storage: at -20°C for 24 months

Ordering information:

Cat.-no	Description	Amount
105-310	Ribonuclease (RNase) Inhibitor	2000 units
105-350	Ribonuclease (RNase) Inhibitor	10000 units

GeneON .. a good decision ..

Contact Phone +49-(0)-621- 5720 864 Fax: +49-(0)-621-5724 462

E-Mail: mailto:info@geneon.net WEB: <http://www.taq-dna.com/> Version: 25.10.2009 AS

Unless specified otherwise, all products of GeneON are sold for research use only.

.. a good decision ..